

MR2888656 (Review) 53A05

Bezkorovaina, L. L. (UKR-ODE-NDM); Vashpanova, T. Yu. (UKR-ODE-NDM)

***A*-deformations of a surface with stationary lengths of *LGT*-lines. (English summary)**

Ukrainian Math. J. **62** (2010), no. 7, 1018–1027.1573-9376

The authors study infinitesimal deformations of a surface that preserve the area element of the surface, which are called *A*-deformations. They define a line of geodesic torsion (*LGT*-line) to be a line on a surface whose direction coincides at every point with a principal direction of geodesic torsion. Then these two notions are combined and *A*-deformations of a surface that preserve the length of lines of geodesic torsion on the surface are examined.

First, a necessary and sufficient condition for the preservation of lengths of *LGT*-lines under an *A*-deformation of a surface is proved. Namely, this condition is the equality

$$2T^{\gamma\delta}c_{\alpha\gamma}g_{\beta\delta} + 2T^{\gamma\delta}c_{\beta\gamma}g_{\alpha\delta} = 2\mu(b_{\alpha\beta} - Hg_{\alpha\beta}),$$

where $\mu(x^1, x^2)$ is a certain function. For a surface with parameter equation $\mathbf{r} = \mathbf{r}(x^1, x^2)$, the $T^{\gamma\delta}$ are deformation tensors on the surface. These are determined by the partial derivatives of the displacement vector $\mathbf{y}(x^1, x^2)$, which in the basis $\mathbf{r}_1, \mathbf{r}_2, \mathbf{n}$ are as follows:

$$\mathbf{y}_i = c_{i\alpha}T^{\alpha\beta}\mathbf{r}_\beta + c_{i\alpha}T^\alpha\mathbf{n}.$$

The $b_{\alpha\beta}$ are the coefficients of the second fundamental form, $c_{i\alpha}$ is the discriminant tensor of the surface ($c_{11} = c_{22} = 0$, $c_{12} = -c_{21} = \sqrt{g}$ and $g = g_{11}g_{22} - g_{12}^2$) and H is the mean curvature of the surface. The $g_{\alpha\beta}$ are the coefficients of the first fundamental form and all the indices take values 1 and 2.

Then the special cases of a trivial *A*-deformation, an *A*-deformation of a minimal surface and an *A*-deformation of a non-flat, non-minimal surface that preserves the lengths of *LGT*-lines are examined. In each of the last two cases, the existence is shown of a nontrivial *A*-deformation with the required property. In the case of a minimal surface, the connection with an *A*-deformation of the surface that preserves the length of asymptotic lines is mentioned.

Also, *A*-deformations with stationary length of *LGT*-lines for surfaces of elliptic and of hyperbolic type are considered. Again, existence results are stated.

In the last section, the deformation tensors of an *A*-deformation that preserves the length of *LGT*-lines are calculated explicitly for an elliptic paraboloid.

Where suitable, the authors point out the similarities and differences between infinitesimal bendings of a surface and *A*-deformations since the latter generalize the former.

Reviewed by *Wendy Goemans*

References

1. I. N. Vekua, *Generalized Analytic Functions* [in Russian], Nauka, Moscow (1988). [MR0977975 \(90a:30135\)](#)

2. P. Vincensini, "Sur les déformations équivalentes infinitésimales des surfaces," *Rev. Univ. Nac. Tucumán A*, **14**, No. 2, 177–188 (1962). [MR0169147 \(29 #6400\)](#)
3. A. D. Aleksandrov, *Convex Polyhedrons* [in Russian], Gostekhteorizdat, Moscow (1950).
4. V. T. Fomenko, "Some results on the theory of infinitesimal bending of surfaces," *Mat. Sb.*, **72** (114), No. 3, 388–411 (1967). [MR0211331 \(35 #2212\)](#)
5. P. G. Kolobov, "On infinitesimal deformations of a surface with preservation of the area," *Uch. Zap. Kabardino-Balkar. Univ., Ser. Mat.*, Issue 30, 65–68 (1966). [MR0225238 \(37 #832\)](#)
6. L. L. Bezkorovaina, "On infinitesimal deformations that preserve the lengths of asymptotic lines," in: *Proceedings of the Scientific Conference of Young Scientists (Natural Sciences)* [in Ukrainian], Odessa (1970), pp. 104–109.
7. L. L. Bezkorovainaya, "Canonical A -deformations that preserve the lengths of curvature lines of a surface," *Mat. Sb.*, **97(139)**, No. 2(6), 163–176 (1975). [MR0394456 \(52 #15257\)](#)
8. N. V. Dermanets, *On Extension of Infinitesimal First-Order Areal Deformations of a Surface of Positive Curvature with Edge to Analytic Ones* [in Russian], Dep. UkrNIINTI, No. 813 Uk-85, Odessa (1985).
9. L. L. Bezkorovaina, *Infinitesimal Areal Deformations and Steady States of an Elastic Shell* [in Ukrainian], Astroprint, Odessa (1999).
10. T. Yu. Vashpanova and L. L. Bezkorovaina, "Geodesic torsion and its extremal values," in: *Proceedings of the Mathematical Scientific Conference of Young Scientists and Students on Partial Differential Equations and Their Applications (Donetsk, December 6–7, 2006)* [in Ukrainian], Donetsk (2006), pp. 28–29.
11. A. V. Bitsadze, *Boundary-Value Problems for Elliptic Equations of the Second Order* [in Russian], Nauka, Moscow (1966). [MR0208152 \(34 #7962\)](#)

Note: This list reflects references listed in the original paper as accurately as possible with no attempt to correct errors.